

Original

Evolution of the antimicrobial resistance of *Pseudomonas aeruginosa* in Spain: Second National Study (2003)

I. Sánchez-Romero,¹ E. Cercenado,² O. Cuevas,² N. García-Escribano,² J. García-Martínez,² E. Bouza,²
and the Spanish Group for the Study of *Pseudomonas aeruginosa*

¹Servicio de Microbiología, Hospital Puerta de Hierro, Madrid, Spain;

²Servicio de Microbiología, Hospital General Universitario Gregorio Marañón, Madrid, Spain.

SUMMARY

The second national prevalence study of *Pseudomonas aeruginosa* has been carried out in Spain. A total of 1250 clinical isolates of *P. aeruginosa* were collected from 127 hospitals in 1 week in 2003 and the resistance data gathered from the isolates was compared with those of the first study in 1998 (1014 isolates from 136 hospitals). Antimicrobial susceptibility testing was performed in both studies in the same laboratory. The most active antimicrobials were piperacillin, piperacillin-tazobactam, and amikacin ($\leq 10\%$ resistant) and resistance to these antimicrobials did not change over the time. The least active were ofloxacin and gentamicin ($\geq 30\%$ resistant). From 1998 to 2003, resistance increased significantly to ciprofloxacin (23% vs. 28%, respectively, $p = 0.015$); ofloxacin (30% vs. 37%, $p = 0.002$); imipenem (14% vs. 18%, $p = 0.017$) and meropenem (8% vs. 13%, $p < 0.001$). Resistance to aztreonam (23%), ceftazidime (16%), cefepime (20%), ticarcillin (13%) and tobramycin (11%) remained stable. Isolates from inpatients were significantly more resistant than those from outpatients to all antimicrobials, with the exception of fluoroquinolones and aminoglycosides ($p < 0.01$). Isolates from outpatients were significantly more resistant to these two groups ($p < 0.05$) than to other antimicrobials. In Spain, from 1998 to 2003, the susceptibility pattern of *P. aeruginosa* to antimicrobial agents has changed. Isolates have become significantly more resistant to fluoroquinolones and carbapenems. However, resistance to beta-lactams and aminoglycosides remains stable.

Key words: *Pseudomonas aeruginosa* - National prevalence study - Antimicrobial resistance

Evolución de la resistencia a los antimicrobianos de *Pseudomonas aeruginosa* en España: Segundo estudio nacional (2003)

RESUMEN

Se ha realizado el segundo estudio nacional de prevalencia de *Pseudomonas aeruginosa* en España. Durante una semana en el año 2003 se recuperaron 1250 aislamientos clínicos de esta bacteria en 127 hospitales españoles y los datos de resistencia se compararon con los obtenidos en el primer estudio realizado en 1998 (1014 aislamientos de 136 hospitales). Las pruebas de sensibilidad se realizaron en el mismo laboratorio en ambos estudios. Los antimicrobianos que se mostraron más activos fueron piperacilina, piperacilina-tazobactam y amikacina

(<10% resistencia), y sus porcentajes de resistencia no cambiaron a lo largo de los dos estudios. Por el contrario, los antimicrobianos menos activos fueron ofloxacino y gentamicina (>30% resistencia). Desde el año 1998 al 2003 se observó un aumento de resistencias significativo en ciprofloxacino (23% vs. 28%, respectivamente, $p=0.015$), ofloxacino (30% vs. 37%, $p=0.002$), imipenem (14% vs. 18%, $p=0.017$) y meropenem (8% vs. 13%, $p<0.001$). Los porcentajes de resistencia a aztreonam (23%), ceftazidima (16%), cefepima (20%), ticarcilina (13%) y tobramicina (11%) permanecieron estables. Los aislamientos de pacientes ingresados fueron significativamente más resistentes que los de los pacientes ambulatorios para todos los antimicrobianos ensayados excepto para las fluoroquinolonas y los aminoglucósidos ($p<0.01$). Los aislamientos procedentes de la comunidad fueron más resistentes significativamente en esos dos grupos ($p<0.05$) que en el resto de los antimicrobianos. En España, los patrones de sensibilidad de *P. aeruginosa* a los antimicrobianos ha cambiado desde 1998 a 2003. Los aislamientos son más resistentes a las fluorquinolonas y los carbapenemes, y se mantienen estables los porcentajes de resistencia para betalactámicos y aminoglucósidos.

Palabras clave: *Pseudomonas aeruginosa* - Estudio de prevalencia nacional - Resistencia antimicrobiana

INTRODUCTION

In 1998, the first national prevalence study of *Pseudomonas aeruginosa* in Spain was carried out; 136 hospitals participated and 1014 isolates were studied (1). The treatment of infections caused by *P. aeruginosa* gives cause for concern, because of its intrinsic resistance to many antimicrobial agents. In addition, *P. aeruginosa* has a high level of ability to acquire resistance via mutations to all relevant antimicrobials and it is frequently seen in severe infections (2-6). The high and increasing rates of local resistance require surveillance studies so that its antimicrobial resistance is known at a given point. However, most studies of the resistance of *P. aeruginosa* to antimicrobial agents are based on the situation in particular institutions or on particular types of patients and clinical syndromes, and thus provide information which is biased. Data about antimicrobial susceptibility without a previous selection are scarce. The results of the second national study, performed in 2003, are reported here and show the evolution of the resistance to antimicrobial agents of *P. aeruginosa* in Spain.

MATERIAL AND METHODS

Participating hospitals

A total of 136 randomly selected Spanish hospitals which represented all types and sizes of public hospitals in Spain and which participated in the first study (1) were invited to take part in the present study, by sending all clinical isolates of *P. aeruginosa* identified during a particular week, without duplication of strains from the same patient and sample, to a reference laboratory. Of the 136 hospitals, 127 participated in this second study. The protocol followed, the reference laboratory, and the identification and susceptibility testing procedures were the same as those used in the first study (1). All isolates were accompanied by a uniform protocol which included the characteristics of the hos-

pital of origin (the number of beds, ward), site of isolation, and acquisition from outpatients (<48 h hospital admission) or inpatients. Information was requested about the population treated by the hospital; the number of admissions per year; the number of samples received by the local microbiology laboratory during the week of the study; and the local susceptibility of *P. aeruginosa* to selected antimicrobial agents. Hospital size was classified into three categories: less than 500 beds; 500-1000 beds; and more than 1000 beds. The ward of isolation was grouped into four categories: high risk of infection (intensive care, units for newborns and immunocompromised patients); surgery; medical units (including pediatrics); and low risk (psychiatry and obstetrics units). The origin of the isolates were classified as: blood; urine; lower respiratory tract; wounds and abscesses; sterile fluids (bile, peritoneal, pleural and cerebrospinal fluid [CSF]); mucosa (conjunctiva, pharyngeal, nasal and urethral); and others.

Identification of the isolates and susceptibility testing

All isolates were identified again at the reference laboratory. Identities and minimum inhibitory concentrations (MICs; mg/l) were determined using an automated microdilution system using MicroScan Neg Combo 1S panels (MicroScan, Baxter Diagnostics, Inc., CA, USA) and following the manufacturer's guidelines. Isolates whose identification was inconclusive were identified again using standard procedures (7) and by the use of the API 20NE system (Biomerieux, Marcy l'Etoile, France). The antimicrobials and concentrations tested were: ticarcillin and piperacillin 16 and 64 mg/l; piperacillin-tazobactam 16/4 mg/l and 64/4 mg/l; ceftazidime, cefepime and aztreonam 1-2 mg/l and 8-16 mg/l; imipenem 1-8 mg/l; meropenem 4-8 mg/l; ciprofloxacin 0.12 and 1-2 mg/l; ofloxacin 0.5 and 2-4 mg/l; gen-

tamicin and tobramycin 4-8 mg/l; and amikacin 8-16 mg/l. Each panel was inoculated with an appropriate dilution of an exponential phase culture of a microorganism. Readings were performed after overnight incubation at 35 °C. *Escherichia coli* ATCC 25922 and *P. aeruginosa* ATCC 27853 were used daily as control strains. Breakpoints were applied following Clinical and Laboratory Standards Institute (CLSI) recommendations (8). MICs (mg/l) in both the intermediate and resistant ranges (CLSI criteria) were considered as nonsusceptible in this study.

Statistical analysis

Susceptibility data were compared by using a Chi-square test (qualitative variables). Statistical analysis was performed with SPSS software version 11.5. In all cases, $p \leq 0.05$ was considered statistically significant. All p-values were calculated as two-sided.

RESULTS

Demographic and identification data

A total of 1270 clinical isolates were received. Of these, 1250 were identified as *P. aeruginosa*. According to the data provided by the participating hospitals, it was estimated that Spanish microbiology laboratories identified 207 *P. aeruginosa* isolates per 100,000 inhabitants of population per year (33 isolates per 1000 hospital admissions

per year). *P. aeruginosa* was recovered in 8% of all the samples received in the microbiology laboratory with bacterial isolates.

The majority of the isolates (43%) were recovered from hospitals with 500-1000 beds. *P. aeruginosa* was recovered in all wards, although the medical wards provided the greatest number of isolates (40%). Nevertheless, considering the smaller number of beds belonging to high-risk areas, the proportion of isolates in these areas was greater. The lower respiratory tract represented the most frequent origin of the isolates (32%), followed by wounds and abscesses (31%) and urine (21%). Thirty per cent of the isolates were recovered from outpatients.

Evolution of resistance of *P. aeruginosa* to antimicrobials

Comparative data regarding antimicrobial resistance in both studies (1998 and 2003) are summarized in Table 1. No one antimicrobial was uniformly active against all *P. aeruginosa* isolates. The most active antimicrobial agents were piperacillin, piperacillin-tazobactam and amikacin ($\leq 10\%$ resistant), and the percentages of resistance to these antimicrobials did not change over time. In contrast, the least active were ofloxacin and gentamicin ($\geq 30\%$ resistant). Resistance rates increased significantly between 1998 and 2003 to ciprofloxacin (23% vs. 28%, respectively, $p=0.015$); ofloxacin (30% vs. 37%, $p=0.002$); imipenem (14% vs. 18%, $p=0.017$) and meropenem (8% vs. 13%, $p<0.001$). Re-

Table 1. In vitro activities of antimicrobial agents against *P. aeruginosa*.

Antimicrobial agent	Year of study						p-value
	1998 (n=1014 isolates)			2003 (n=1250 isolates)			
	Range (mg/l)	MIC ₉₀ (mg/l)	%R*	Range (mg/l)	MIC ₉₀ (mg/l)	%R*	
Amikacin	≤8->16	16	9	≤8->16	16	8	NS
Aztreonam	≤1->16	16	23	≤1->16	16	23	NS
Cefepime	≤1->16	16	17	≤1->16	16	20	NS
Ceftazidime	≤1->16	16	15	≤1->16	16	16	NS
Ciprofloxacin	≤0.12->2	>2	23	≤0.12->2	>2	28	0.015
Gentamicin	≤4->8	>8	31	≤4->8	>8	30	NS
Imipenem	≤1->8	8	14	≤1->8	>8	18	0.017
Meropenem	≤4->8	4	8	≤4->8	8	13	<0.001
Ofloxacin	≤0.5->4	>4	30	≤0.5->4	>4	37	0.002
Piperacillin	≤16->64	64	10	≤16->64	64	10	NS
Piperacillin-tazobactam	≤16->64	64	7	≤16->64	64	7	NS
Ticarcillin	≤16->64	>64	13	≤16->64	>64	13	NS
Tobramycin	≤4->8	4	10	≤4->8	>8	11	NS

MIC: minimum inhibitory concentration.

*Percentage of resistance, includes all nonsusceptible (intermediate and resistant) isolates. NS: statistically nonsignificant.

Table 2. Resistance* of *P. aeruginosa* isolates from inpatients and outpatients.

Antimicrobial agent	Inpatients (n=875)	Outpatients (n=375)	p-value
Amikacin	8	7	NS
Aztreonam	26	17	0.002
Cefepime	23	15	0.004
Ceftazidime	20	9	<0.001
Ciprofloxacin	29	26	NS
Gentamicin	32	28	NS
Imipenem	24	9	<0.001
Meropenem	18	5	<0.001
Ofloxacin	39	34	NS
Piperacillin	13	6	<0.001
Piperacillin-tazobactam	10	2	<0.001
Ticarcillin	16	8	<0.001
Tobramycin	13	9	NS

*Percentage of resistance, includes all nonsusceptible (intermediate and resistant) isolates. NS: statistically nonsignificant.

sistance to aztreonam (23%), ceftazidime (16%), cefepime (20%), ticarcillin (13%), and tobramycin (11%) remained stable. Isolates from inpatients were significantly more resistant than those from outpatients to all antimicrobials, with the exception of fluoroquinolones and aminoglycosides (Table 2). Isolates from outpatients were significantly more resistant to these two groups of antimicrobials than

to others ($p < 0.05$). Resistance to aminoglycosides, with the exception of amikacin, and to carbapenems were significantly higher in hospitals with more than 1,000 beds ($p < 0.05$). Isolates from high-risk areas were more resistant to carbapenems and beta-lactams than those from other wards ($p < 0.01$). Isolates from catheter tips were significantly more resistant to all antimicrobials tested ($p < 0.01$) than those from other samples; only 17 isolates came from this origin. In contrast, isolates from urine were the most susceptible to all antimicrobials with the exception of fluoroquinolones: a significant increase in the resistance to ciprofloxacin of urine isolates was observed in comparison with the results obtained in the first study (28% in 1998 vs. 38% in 2003, $p = 0.020$). For outpatient isolates, urine was the most common site of isolation (31%); this was significantly more common than lower respiratory tract ($p < 0.05$). In contrast, in strains from inpatients lower respiratory tract was the most common site of isolation and it was significantly more common than urine ($p < 0.05$). The cross-resistance of *P. aeruginosa* isolates to antimicrobial agents is summarized in Table 3. The majority (93%) of meropenem-resistant isolates were also resistant to imipenem; however, 33% of imipenem-resistant isolates were susceptible to meropenem. About half of the imipenem-resistant isolates were susceptible to ceftazidime, cefepime, aztreonam and ciprofloxacin, and 75% were sus-

Table 3. Crossresistance of *P. aeruginosa* isolates.

Drug(s) to which isolates were resistant	% Resistance* to												
	AK	AZT	CPE	CAZ	CIP	GN	IMP	MER	OFL	PI	P/T	TI	TO
AK		45	59	47	66	100	45	37	78	36	25	37	58
AZT	15		60	51	53	50	41	38	72	37	27	51	24
CPE	36	92		92	56	77	66	62	71	74	68	83	34
CAZ	28	94	95		56	61	58	53	67	81	61	78	32
CIP	28	43	43	33		62	34	27	99	21	15	29	35
GN	25	27	46	32	57		36	29	67	22	16	27	36
IMP	19	52	53	47	54	61		67	67	30	25	34	35
MER	22	67	68	61	60	68	93		76	39	34	46	43
OFL	16	45	40	29	76	56	33	27		18	13	28	28
PI	28	83	92	89	60	67	52	50	67		64	78	42
P/T	29	90	98	93	61	73	65	64	70	96		92	39
TI	23	91	83	69	64	64	48	47	81	62	48		36
TO	40	48	67	59	87	97	55	49	91	38	23	40	

*Includes all nonsusceptible (intermediate and resistant) isolates.

AK: amikacin; AZT: aztreonam; CPE: cefepime; CAZ: ceftazidime; CIP: ciprofloxacin; GN: gentamicin; IMP: imipenem; MER: meropenem; OFL: ofloxacin; P/T: piperacillin-tazobactam; PI: piperacillin; TI: ticarcillin; TO: tobramycin.

ceptible to piperacillin-tazobactam. About 40% of the ceftazidime-resistant isolates were susceptible to carbapenems, piperacillin-tazobactam and ciprofloxacin. All amikacin nonsusceptible isolates were resistant to gentamicin, but 42% were susceptible to tobramycin ($MIC \leq 4$ mg/l). Around 75% of the gentamicin-resistant isolates were susceptible to amikacin and tobramycin. Approximately two-thirds of the ciprofloxacin-resistant isolates were susceptible to ceftazidime, piperacillin-tazobactam, carbapenems, tobramycin and amikacin.

DISCUSSION

This study demonstrates that important changes have occurred in the susceptibility pattern of *P. aeruginosa* to antimicrobial agents in a short period of time in Spain. Isolates have become significantly more resistant to fluoroquinolones and carbapenems, however, resistance to beta-lactams and aminoglycosides has remained stable. Infections due to *P. aeruginosa* are often difficult to treat due to its virulence and to the relatively limited choice of effective antimicrobial agents (9, 10). In recent years, several studies have reported an increasing resistance of *P. aeruginosa* to different antimicrobial agents (11-15). Most studies, however, report the resistance of *P. aeruginosa* to a particular class of antimicrobials (16-18), in special units or diseases (19-21), or in special types of patients (22, 23). Our study is not limited to specific aspects and includes all isolates without *a priori* selection. In a multicenter study performed in Belgium and Luxembourg in 1999 and in which 40 hospitals participated, the percentages of resistance of *P. aeruginosa* to meropenem (9.5%) and to ciprofloxacin (24%) were similar to those of the present study (24). Another study in Italy showed similar percentages of resistance to ciprofloxacin (29% in 1998 and 28% in 2002) (25). In a multicenter study in the United States, resistance to fluoroquinolones was 29% in 1999 and 36% in 2001 (26). In our study, a high percentage of *P. aeruginosa* isolates were obtained from outpatients, of which 26% were resistant to ciprofloxacin and 34% to ofloxacin. The low percentages of susceptibility to fluoroquinolones may reflect the high use of these antimicrobials in the community. The present study demonstrates that beta-lactams, despite having been in use for a longer time, have higher *in vitro* activity than the fluoroquinolones against *P. aeruginosa*; resistance to beta-lactams has not increased in comparison with the first study performed in 1998. It is noteworthy that the percentages of resistance to ceftazidime (16%) and piperacillin-tazobactam (7%) have remained stable. Resistance to cefepime showed a tendency to increase (16% in 1998 vs. 20% in 2003), but this did not reach statistical significance. The SENTRY Antimicrobial

Surveillance Program (1998-2003) performed in U.S. medical centers reported 11.6% of ceftazidime-resistant *P. aeruginosa* (27), while another SENTRY study (1998-2004) reported 16.9% of isolates with resistance to ceftazidime (28). Other studies have demonstrated similar values of resistance to carbapenems, ciprofloxacin and aminoglycosides to those found in the present study (29-31). Isolates from inpatients were significantly more resistant to all antimicrobials with the exception of fluoroquinolones and aminoglycosides, but isolates from outpatients presented high levels of resistance to these antimicrobials as well. We observed significantly higher percentages of resistance to carbapenems and aminoglycosides in hospitals with more than 1000 beds, but found no relevant data in the literature. Significantly higher percentages of resistance to carbapenems and beta-lactams were also observed in isolates from high-risk areas and that may reflect the higher use of these antimicrobial agents in the intensive care units (ICUs). The high rate of resistance to all antimicrobials in isolates from catheter tips may be due to the scarce number of isolates with this origin in the present study. On the other hand, isolates from urine were the most susceptible to all antimicrobials tested, with the exception of fluoroquinolones. These data are in accordance with previous publications (32). Cross-resistance data indicate that a high number of isolates probably have resistance due to either a combination of multiple unrelated mechanisms of resistance or the involvement of efflux pump overexpression in association with fluoroquinolone resistance.

CONCLUSION

The intrinsic resistance of *P. aeruginosa* to many antimicrobial agents and the high ability of this microorganism to acquire new mechanisms of resistance make its management problematic in the hospital setting. However, in our study, there are still several antimicrobials (carbapenems, cefepime, ceftazidime, tobramycin and amikacin) that have good activity against *P. aeruginosa*, with piperacillin-tazobactam being the most active agent. Care must be taken in the overuse of carbapenems as this may lead to increasing resistance to these drugs, as observed in the present study.

In summary, in a very short period of time significant changes have been observed in the resistance of *P. aeruginosa* to antimicrobial agents in Spain that are not attributable to a selected institution or to a selected type of patient. Periodical surveillance studies are essential to determine the current susceptibility of *P. aeruginosa* to different antimicrobial agents and to provide an overview of the situation in a given country.

ACKNOWLEDGMENTS

This work was supported in part by "Red Española de Investigación en Patología Infecciosa" (REIPI - ISCIII - C03/14).

The members of the Spanish Group for the Study of *Pseudomonas aeruginosa* and the staff of the microbiology services of all participating hospitals are as follows: M. Rodríguez-Jove, A. Álvarez, Hospital General Juan Cardona (El Ferrol-La Coruña); J.A. Agulla, M. Rodríguez-Mayo, Complejo Hospitalario Arquitecto Marcide-Novoa Santos (El Ferrol-La Coruña); B. Regueiro, F. Pardo, Hospital de Conxo (Santiago de Compostela-La Coruña); P. Alonso, A. Coira, Complejo Hospitalario Xeral-Calde (Lugo); A. Tinajas, Complejo Hospitalario Cristal Piñor (Orense); V. Pulian, M. García-Campello, Complejo Hospitalario de Pontevedra (Pontevedra); T. González-Blanco, I. Otero, Complejo Hospitalario Xeral-Cies (Vigo-Pontevedra); J. Torres, F.J. Vasallo, Hospital do Meixoeiro (Vigo-Pontevedra); J. Sevillano, I. Rodríguez-Conde, Policlínico POVISA (Vigo-Pontevedra); F. Vázquez, M. Aranaz, Hospital Monte Naranco (Oviedo-Asturias); F. Méndez, M. Lantero, Hospital Central de Asturias (Oviedo-Asturias); E. Hidalgo, Hospital de Jove (Gijón-Asturias); G. Viejo, M.D. Miguel, Hospital de Cabueñas (Gijón-Asturias); P. Prendes, J. Rodríguez-Álvarez, Hospital San Agustín (Avelés-Asturias); R. Cimadevilla, Hospital Comarcal de Jario (Asturias); A. Torreblanca, Hospital Carmen y Severo Ochoa (Cangas de Narcea-Asturias); L. Martínez, J. Calvo, Hospital Marqués de Valdecilla (Santander); L.F. Colomo, P. Mellado, Hospital Comarcal de Laredo (Cantabria); R. Cisterna, K. Ibarra, Hospital de Basurto (Bilbao-Vizcaya); F. Calvo, Hospital de Santa Marina (Bilbao-Vizcaya); I. Marzana, G. Martín-Saco, Hospital San Eloy (Baracaldo-Vizcaya); M. Alkorta, J. Barrón, Hospital de Cruces (Baracaldo-Vizcaya); M.J. López-Goikoetxea, Hospital de Galdakao (Vizcaya); L. Michaus, M. de Pablos, Hospital Txagorritxu (Vitoria); A. Labora, A. Canut, Hospital Santiago Apóstol (Vitoria); E. Pérez-Trallero, J.M. García-Arenzana, Hospital de Donostia (Guipúzcoa); J.A. Jiménez-Alfaro, A. Jauregui, Policlínica Guipúzcoa (San Sebastián-Guipúzcoa); L. Torroba, Hospital Virgen del Camino (Pamplona); C. Pina, A. Fontaneda, Hospital San Juan de Dios (Pamplona); I. Dorronsoro, J.J. García-Irure, Hospital de Navarra (Pamplona); R. Díaz-García, J. Leiva, Clínica Universitaria de Navarra (Pamplona); M.J. Gastañares, I. Olarte, Complejo Hospitalario San Millan-San Pedro (Logroño); M.T. Jiménez-Anta, F. Marco, Hospital Clinic i Provincial (Barcelona); P. Coll, B. Mirelis, Hospital de la Santa Creu i Sant Pau (Barcelona); R. Martín, F. Tubau, Hospital Princeps d'Espanya (Hospitalet Llobregat-Barcelona); G. Prats, N. Larrosa, Ciutat Sanitaria Vall D'Hebron (Barcelona); M. Salvadó, Hospital del Mar (Hospitalet Llobregat-Barcelona); D. Fontanals, D. Mariscal, Consorcio Hospitalario del Parc Taulí (Sabadell-Barcelona); J. Lite, Mutua de Tarrasa (Barcelona); V. Ausina, L. Matas, Hospital Universitari Germans Trias i Pujol (Badalona-Barcelona); F. Corcoy, R. Angrill, Residencia Sant Camil (Sant Pere de Ribes-Barcelona); M.L. Urcula, Hospital Provincial de Santa Caterina (Gerona); J. Batlle, M. Motje, Hospital Universitario Doctor Josep Trueta (Gerona); A. García-Busto, R. Moreno, Hospital General (Castellón); M. Canós, B. Vila, Hospital La Plana (Villareal-Castellón); M. Gobernado, J.L. López-Hontangas, Hospital Universitario La Fe (Valencia); J. García-Lomas, D. Navarro, Hospital Clínico (Valencia); A. Lloret, M. Bosque, Hospital Arnau Vilanova (Valencia); J. Maiquez, E. Aznar, Instituto Valenciano de Oncología (Valencia); J.M. Nogueira, A. Morales, Hospital Dr. Peset (Valencia); M.R. Llucian, Hospital General Universitario (Valencia); J.M. García-Aguayo, Hospital General de Requena, (Valencia); M.C. Alonso, L. Andreu, Hospital Francesc Borja (Gandía-Valencia); D. González-Granda, Hospital Lluís Alcanyis (Játiva-Valencia); J.L. Hernández, Hospital Militar Vázquez Bernabeu (Quart Poblet-Valencia); J. Prat, R. Escoms,

Hospital Sagunto (Valencia); S. Giner, Hospital Dr. Moliner (Serra-Valencia); A. Yagüe, N. Gonzalo, Hospital de la Vega Baja (San Bartolomé-Alicante); G. Royo, L. Cebrian, Hospital General Universitario (Elche-Alicante); A. Altuna, Hospital General Universitario (Murcia); M. Segovia, A. Menasalvas, Hospital Morales Meseguer (Murcia); J. Piqueras, L. Martínez, J.M. Sicilia, Hospital General Santa María del Rosell (Cartagena-Murcia); J. Ruiz, V. Vilar, Hospital Universitario Virgen de la Arrixaca (El Palmar-Murcia); J.L. Pérez, N. Borrell, A. Oliver, Hospital Son Dureta (Palma de Mallorca); J. Sánchez-Gómez, Hospital Can Misses (Ibiza); A. Gutiérrez, A. García-Perea, Hospital La Paz (Madrid); J.J. Rodríguez-Otero, F. Chaves, Hospital Universitario Doce de Octubre (Madrid); M. Menéndez-Rivas, Hospital del Niño Jesús (Madrid); F. Hervas, P. Sánchez, Hospital Militar Gómez Ulla (Madrid); J.J. Picazo, A. San Pedro, Hospital Clínico San Carlos (Madrid); F. Baquero, R. Cantón, Hospital Ramón y Cajal (Madrid); M. A. Blanco, C. Pazos, Hospital Santa Cristina (Madrid); D. Dámaso, Hospital Puerta de Hierro (Madrid); M. López-Brea, T. Alarcón, Hospital Universitario de la Princesa (Madrid); R. Cortés, V. Portús, Hospital de la Cruz Roja -San José y Santa Adela (Madrid); A. Urmeneta, Hospital de Cantoblanco (Madrid); M. Beltrán, P. Gómez, Hospital Príncipe de Asturias (Alcalá de Henares-Madrid); J.L. Gómez, J. Tamayo, Hospital General (Móstoles-Madrid); I. Wilhelmi-Cal, Hospital Severo Ochoa (Leganés-Madrid); M. Sánchez, J. Cacho, Hospital Universitario (Getafe-Madrid); E. Rollán, T. Pérez, Hospital Militar (Zaragoza); C. Miñiana, Hospital N. Señora de Gracia (Zaragoza); M.J. Revillo, Hospital Miguel Servet (Zaragoza); C. Rubio, R. Escartín, Hospital Clínico Universitario Lozano Blesa, (Zaragoza); M. Ferrero, Hospital General San Jorge (Huesca); P. Chocarro, Hospital General Obispo Polanco (Teruel); C. Navarro, T. Nebreda, Hospital de Alcañiz, (Teruel); L. Díaz, S. Brea, Hospital Virgen De La Salud (Toledo); A.M. Leturia, Hospital Nacional de Paraplégicos (Toledo); J.C. González-Rodríguez, I. Barba, Hospital Nuestra Señora del Carmen (Ciudad Real); M.D. Romero, F. Mora, Hospitales Alarcos y Carmen (Ciudad Real); J. Bisquert, M.T. Pérez, Hospital General Universitario (Guadalajara); D. Crespo, E. Escribano, P. Robles, Complejo Hospitalario (Albacete); F. Cachón, Hospital de León (León); C. Fuster, Hospital del Bierzo (Ponferrada-León); M.F. Brezmes, L. López-Urrutia, Hospital Virgen de La Concha, (Zamora); J.A. García-Rodríguez, J.E. García-Sánchez, Hospital Virgen de La Vega (Salamanca); S. García-Carbajosa, P. Carrero, Hospital General (Segovia); I. Pozas, E. Ojeda, Mejías, Hospital General Yagüe (Burgos); M.D. Badía, C. Lizondo, Hospital Provincial Divino Vallés (Burgos); C. Gimeno, Hospital Comarcal Santiago Apóstol (Miranda de Ebro-Burgos); R. Ibáñez, A. Gómez, Hospital Ntra. Sra. de Sonsoles (Ávila); A. Campos, F. Merino, Hospital General (Soria); M. A. García-Castro, E. Álvarez, Hospital General Río Carrión (Palencia); A. Rodríguez-Torres, M. A. Bratos, Hospital Universitario (Valladolid); R. Iñiguez, Hospital Nuestra Señora de La Montaña (Cáceres); P. Teno, Complejo Hospitalario San Pedro de Alcántara (Cáceres); J. Blanco, E. Garduño, Hospital Universitario Infanta Cristina (Badajoz); J. García-Herruzo, A. Saldarreaga, M. Martín, Hospital Universitario Puerta Del Mar (Cádiz); I. Ruiz, Hospital Punta de Europa (Algeciras-Cádiz); L. Calbo, J.L. de Francisco, Hospital de Jerez de La Frontera (Cádiz); A. Sánchez-Porto, Hospital La Línea de La Concepción (Cádiz); M. Casal, A. Ibarra, Hospital Universitario Reina Sofía (Córdoba); M. Rosa-Fraile, C. Miranda, Hospital Virgen de Las Nieves (Granada); P. Manchado, J. Porras, Hospital Carlos Haya (Málaga); I. Cuesta, C. Carazo, Hospital Universitario (Jaén); J.M. Saavedra, L. Pascual, Hospital Juan Ramón Jiménez (Huelva); C. García-Iglesias, Hospital San Lázaro (Sevilla); M. Sánchez, M. Chavez, Hospital San Juan de Dios (Sevilla); E. Perea, E. Ramírez, Y. Guerrero, Hospital Virgen Macarena (Sevilla); J. Aznar, L. Merino, Hospital Virgen del Rocío (Sevilla); E. Martín, Hos-

pital Nuestra Señora de Valme (Dos Hermanas-Sevilla); J. López-Barba, J. Díaz, Hospital Ingesa de Ceuta (Ceuta); M. Galán, Hospital Comarcal (Melilla); J.A. Tur, Hospital Militar Pages (Melilla); N. Batista, A. Moreno, Hospital Nuestra Señora de La Candelaria (Santa Cruz de Tenerife); A. Sierra, M. Cuervo, Hospital Universitario de Canarias (La Laguna-Santa Cruz de Tenerife); R. M. Gallardo, Hospital General de La Palma (Santa Cruz de La Palma-Santa Cruz de Tenerife); A.M. Martín, M. Bollaños, Hospital Insular de Gran Canaria (Las Palmas de Gran Canaria); A. Fleites, M. J. Santos, Hospital General de Asturias H.U.C.A. (Oviedo); G. Esteban, B. Fernández, Hospital Santa María Nai (Ourense); A. Guerrero, M. Cuenca, P. Ramos, Hospital de la Rivera (Alzira).

Correspondence: Emilia Cercenado, Servicio de Microbiología, Hospital General Universitario Gregorio Marañón, C/ Dr. Esquerdo nº 46, 28007 Madrid, Spain. Tel.: +34-91-586-8459; Fax: +34-91-504-4906; E-mail: ecercenado@terra.es

REFERENCES

- Bouza, E., García-Garrote, F., Cercenado, E., Marín, M., Díaz, M.S. *Pseudomonas aeruginosa: A survey of resistance in 136 hospitals in Spain. The Spanish Pseudomonas aeruginosa Study Group.* Antimicrob Agents Chemother 1999; 43: 981-982.
- Cavallo, J.D., Plesiat, P., Couetdic, G., Leblanc, F., Fabre, R. *Mechanisms of beta-lactam resistance in Pseudomonas aeruginosa: Prevalence of OprM-overproducing strains in a French multicentre study (1997).* J Antimicrob Chemother 2002; 50: 1039-1043.
- Hooper, D.C. *Emerging mechanisms of fluoroquinolone resistance.* Emerg Infect Dis 2001; 7: 337-341.
- Juan, C., Macia, M.D., Gutiérrez, O. et al. *Molecular mechanisms of beta-lactam resistance mediated by AmpC hyperproduction in Pseudomonas aeruginosa clinical strains.* Antimicrob Agents Chemother 2005; 49: 4733-4738.
- Okamoto, K., Gotoh, N., Nishino, T. *Pseudomonas aeruginosa reveals high intrinsic resistance to penem antibiotics: Penem resistance mechanisms and their interplay.* Antimicrob Agents Chemother 2001; 45: 1964-1971.
- Pai, H., Kim, J., Lee, J.H., Choe, K.W., Gotoh, N. *Carbapenem resistance mechanisms in Pseudomonas aeruginosa clinical isolates.* Antimicrob Agents Chemother 2001; 45: 480-484.
- Kiska, D.L., Gilligan, P.H. *Pseudomonas.* In: Murray, P.R. et al. (Eds.). *Manual of Clinical Microbiology*, 7th ed. ASM Press, Washington DC 1999; pp. 517-525.
- CLSI. *Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing*, Vol. 26, No. 3, 16th informational supplement. M100-S16. Clinical and Laboratory Standards Institute, Wayne, PA 2006.
- Ferrara, A.M. *Potentially multidrug-resistant non-fermentative Gram-negative pathogens causing nosocomial pneumonia.* Int J Antimicrob Agents 2006; 27: 183-195.
- Ferreira, A.C., Gobara, S., Costa, S.E. et al. *Emergence of resistance in Pseudomonas aeruginosa and Acinetobacter species after the use of antimicrobials for burned patients.* Infect Control Hosp Epidemiol 2004; 25: 868-872.
- McGowan, J.E., Jr. *Resistance in nonfermenting gram-negative bacteria: Multidrug resistance to the maximum.* Am J Med 2006; 119: S29-36.
- Lister, P.D., Wolter, D.J., Wickman, P.A., Reisbig, M.D. *Levofloxacin/imipenem prevents the emergence of high-level resistance among Pseudomonas aeruginosa strains already lacking susceptibility to one or both drugs.* J Antimicrob Chemother 2006; 57: 999-1003.
- Hocquet, D., Nordmann, P., El Garch, F., Cabanne, L., Plesiat, P. *Involvement of the MexXY-OprM efflux system in emergence of cefepime resistance in clinical strains of Pseudomonas aeruginosa.* Antimicrob Agents Chemother 2006; 50: 1347-1351.
- Deplano, A., Denis, O., Poirel, L. et al. *Molecular characterization of an epidemic clone of panantibiotic-resistant Pseudomonas aeruginosa.* J Clin Microbiol 2005; 43: 1198-1204.
- Walsh, F.M., Amyes, S.G. *Microbiology and drug resistance mechanisms of fully resistant pathogens.* Curr Opin Microbiol 2004; 7: 439-444.
- Husson, M.O., Richet, H., Aubert, A. et al. *In vitro comparative activity of meropenem with 15 other antimicrobial agents against 1798 Pseudomonas aeruginosa isolates in a French multicenter study.* Clin Microbiol Infect 1999; 5: 499-503.
- Quentin, C., Tessier, F., Abinaser, A. et al. *Comparative in vitro activity of cefepime: Multicenter study in Aquitaine.* Pathol Biol (Paris) 1997; 45: 363-370.
- Dubrous, P., Cavallo, J.D., Buisson, Y. *Sensitivity to fosfomycin of multiresistant serotype 012 Pseudomonas aeruginosa. Multicenter study.* Pathol Biol (Paris) 1997; 45: 472-478.
- Eigen, H., Rosenstein, B.J., Fitzsimmons, S., Schidlow, D.V. *A multicenter study of alternate-day prednisone therapy in patients with cystic fibrosis. Cystic Fibrosis Foundation Prednisone Trial Group.* J Pediatr 1995; 126: 515-523.
- García-Rodríguez, J.A., Trujillano Martín, I., Baquero, F. et al. *In vitro activity of fosfomycin trometamol against pathogens from urinary tract infections: A Spanish multicenter study.* J Chemother 1997; 9: 394-402.
- Gehanno, P. *Multicenter study of the efficacy and safety of oral ciprofloxacin in the treatment of chronic suppurative otitis media in adults. The French Study Group.* Otolaryngol Head Neck Surg 1997; 117: 83-90.
- De Medeiros, E.A. *Treatment of pneumonia in hospitalized patients: Results of a multicenter study using a fourth-generation cephalosporin (cefepime).* Rev Assoc Med Bras 1999; 45: 2-8.
- Donati, L., Periti, P., Andreassi, A. et al. *Increased burn patient survival with once-a-day high dose teicoplanin and netilmicin. An Italian multicenter study.* J Chemother 1998; 10: 47-57.
- Van Eldere, J. *Multicentre surveillance of Pseudomonas aeruginosa susceptibility patterns in nosocomial infections.* J Antimicrob Chemother 2003; 51: 347-352.
- De Vecchi, E., Drago, L., Nicola, L. et al. *Resistance of Pseudomonas aeruginosa to ciprofloxacin and levofloxacin: 1998-2002.* Infez Med 2003; 11: 196-200.
- Polk, R.E., Johnson, C.K., McClish, D., Wenzel, R.P., Edmond, M.B. *Predicting hospital rates of fluoroquinolone-resistant Pseudomonas aeruginosa from fluoroquinolone use in US hospitals and their surrounding communities.* Clin Infect Dis 2004; 39: 497-503.
- Sader, H.S., Fritsche, T.R., Jones, R.N. *Potency and spectrum trends for cefepime tested against 65746 clinical bacterial isolates collected in North American medical centers: Results from the SENTRY Antimicrobial Surveillance Program (1998-2003).* Diagn Microbiol Infect Dis 2005; 52: 265-273.
- Pfaller, M.A., Sader, H.S., Fritsche, T.R., Jones, R.N. *Antimicrobial activity of cefepime tested against ceftazidime-resistant Gram-negative clinical strains from North American Hospitals: Report from the*

- SENTRY Antimicrobial Surveillance Program (1998-2004)*. *Diagn Microbiol Infect Dis* 2006; 56: 63-68.
29. Picazo, J.J., Betriu, C., Rodríguez-Avial, I., Culebras, E., Gómez, M. *Surveillance of antimicrobial resistance: VIRA study 2004*. *Enferm Infecc Microbiol Clin* 2004; 22: 517-525.
 30. Astal, Z. *Susceptibility patterns in Pseudomonas aeruginosa causing nosocomial infections*. *J Chemother* 2004; 16: 264-268.
 31. Gomila Sard, B., Pardo Serrano, F.J., Moreno Muñoz, R., Celades Porcar, E., García Del Busto Remón, A. *Antimicrobial susceptibility of Pseudomonas aeruginosa clinical isolates in Castellon, Spain*. *Rev Esp Quimioterap* 2006; 19: 60-64 .
 32. Muratani, T., Matsumoto, T. *Bacterial resistance to antimicrobials in urinary isolates*. *Int J Antimicrob Agents* 2004; 24(Suppl. 1): S28-31.