

Brief report

Pilar Escribano^{1,2}
María Jesús Pérez-Granda^{1,2,3}
Roberto Alonso^{1,2,3,4}
Pilar Catalán^{1,2,3}
Luis Alcalá^{1,2,3}
José Antonio Serra-Rexarch^{2,4,5,6}
Laura Osuna^{2,5}
Alejandro Fernández^{2,5}
Aurora Conti^{2,5}
Alberto Castellanos^{2,5}
Jesús Guinea^{1,2,3}
Patricia Muñoz^{1,2,3,4}
Emilio Bouza^{1,2,3,4}

High incidence of COVID-19 at nursing homes in Madrid, Spain, despite preventive measures

¹Clinical Microbiology and Infectious Diseases, Hospital General Universitario Gregorio Marañón, Madrid, Spain

²Instituto de Investigación Sanitaria Gregorio Marañón, Madrid, Spain

³CIBER Enfermedades Respiratorias-CIBERES (CB06/06/0058), Madrid, Spain

⁴Medicine Department, Faculty of Medicine, Universidad Complutense de Madrid, Madrid, Spain

⁵Department of Geriatrics, Hospital General Universitario Gregorio Marañón, Madrid, Spain

⁶CIBER de Fragilidad y Envejecimiento Saludable (CIBERFES), Madrid, Spain

Article history

Received: 22 January 2022; Revision Requested: 19 February 2022; Revision Received: 22 February 2022;

Accepted: 17 March 2022; Published: 11 April 2022

ABSTRACT

Objective. To assess the impact of COVID-19 at nine nursing homes in Madrid, Spain, during the first wave of COVID-19 infection and lockdown period when preventive measures were taken to avoid transmission among residents.

Methods. Nine hundred forty-two residents and 846 staff members from nine nursing homes participated in the study (April 18 to June 20, 2020). All participants were tested for SARS-CoV-2 in the nasopharynx by PCR and for IgG antibodies detection. Microbiological status at sampling was defined as active infection (positive PCR ± presence of antibodies), past infection (negative PCR + presence of antibodies), or naïve participants (negative PCR + absence of antibodies).

Results. Laboratory results helped classify the residents as having active infection (n=224; 23.8%), past infection (n=462; 49.1%), or being naïve (n=256; 27.1%); staff members were actively infected (n=127; 15.1%), had had a past infection (n=290; 34.2%), or were naïve (n=429; 50.7%). Overall, the percentage of participants with COVID-19 was significantly higher in residents than in staff members (72.8% vs 49.2%; $P=0.001$). The clinical situation of residents vs staff at sampling was as follows: acute manifestations compatible with COVID-19 (7.3% vs 3.9%; $P<0.01$) and no manifestations of infection (92.7% vs 96.0%; $P<0.01$). A large proportion of both asymptomatic and symptomatic residents (69.4% vs 86.6%; $P=0.015$) had positive PCR results (mostly alongside positive IgG determinations).

Conclusions. COVID-19 affects 75% of the residents in nursing homes in Madrid. The high impact in these settings,

despite the strict restrictions adopted during the lockdown, demonstrates the ability of SARS-CoV-2 to cause outbreaks.

Keywords: COVID-19, nursing home, PCR, antibody, IgG

Elevada incidencia de COVID-19 en residencias de ancianos en Madrid, España, a pesar de las medidas de prevención

RESUMEN

Objetivo. Evaluar el impacto de la COVID-19 en nueve residencias de ancianos en Madrid (España) durante la primera ola de infección y el período de confinamiento, cuando se tomaron medidas preventivas para evitar la transmisión en estos centros.

Métodos. Se tomaron muestras de 942 residentes y 846 miembros del personal de nueve residencias de ancianos (del 18 de abril al 20 de junio de 2020). A todos los participantes se les realizó una prueba PCR en muestra nasofaríngea, y detección de anticuerpos IgG en sangre. El estado microbiológico en el momento del muestreo se definió como infección activa (PCR positiva ± presencia de anticuerpos), infección pasada (PCR negativa + presencia de anticuerpos) o sin infección (PCR negativa + ausencia de anticuerpos).

Resultados. Los residentes tuvieron infección activa (n=224; 23,8%), infección pasada (n=462; 49,1%) o no tuvieron infección (n=256; 27,1%); el personal presentó infección activa (n=127; 15,1%), infección pasada (n=290; 34,2%) o no tuvieron infección (n=429; 50,7%). En general, el porcentaje de participantes con COVID-19 fue significativamente mayor entre los residentes que entre los miembros del personal (72,8% vs 49,2%; $P=0,001$). La situación clínica de los residentes vs el personal en el momento del muestreo fue: manifestaciones agudas compatibles con COVID-19 (7,3% vs 3,9%; $P<0,01$) y sin manifestaciones de infección (92,7% vs 96,0%; $P<0,01$). Una elevada proporción de residentes tanto asintomáticos co-

Correspondence:

Pilar Escribano

Servicio de Microbiología Clínica y Enfermedades Infecciosas, Hospital General Universitario Gregorio Marañón

C/ Dr. Esquerdo, 46, 28007 Madrid, Spain

Phone +34 91 5867163; Fax +34 91 3721721

E-mail: pilar.escribano.martos@gmail.com

mo sintomáticos (69,4% vs 86,6%; $P=0,015$) tuvieron resultados positivos de PCR (principalmente junto con determinaciones positivas de IgG).

Conclusiones. La COVID-19 afectó al 75% de los ancianos de las residencias de mayores de Madrid. El alto impacto en este entorno, a pesar de las estrictas restricciones adoptadas durante el confinamiento, demuestra la capacidad del SARS-CoV-2 para provocar brotes.

Palabras clave: COVID-19, residencias de ancianos, PCR, anticuerpos, IgG

INTRODUCTION

Spain has been one of the hardest hit countries by the COVID-19 pandemic [1] and the elderly population has been the most affected [2]. COVID-19 disproportionately affected residents and care workers at nursing homes during the first wave of infections that started late-February 2020 and lasted for three months [3]. The exponential increase of cases overwhelmed the Spanish healthcare system and, during the lockdown (from March 15 to June 21), family visits to nursing homes were forbidden. Despite the measures, SARS-CoV-2 kept spreading, causing multiple outbreaks with high-associated mortality rate [4].

The real impact of COVID-19 in nursing homes may be underestimated given the high number of residents or staff members who were asymptomatic and/or the lack of proper testing at the beginning of the pandemic. Epidemiological studies combining both clinical and laboratory data can help to better understand the extent and spreading of SARS-CoV-2 infections in critical settings during the pandemic. The combination of PCR testing and antibody determination helped us to describe that up to 92% and 48% of residents and staff members, respectively, were infected with COVID-19 in a nursing home in Madrid [5]. We here assessed the impact of COVID-19 at nursing homes in Madrid, Spain, during the first wave of COVID-19 infection, when first strict lockdown period and initial preventive measures were implemented.

MATERIAL AND METHODS

All surviving residents and health care workers who consented to participate in the present study were included. A total of 1,788 residents and staff members from nine nursing homes located in Madrid, Spain, participated in the study between April 18 and June 20, 2020. Subjects were assessed with both a nasopharynx PCR sample for PCR testing (TaqMan™ 2019-nCoV assay, Applied Biosystems, Pleasanton, CA, USA). Positive PCR results were defined as simultaneous amplification of ORF, S and N genes with Ct values lower than 37. Detection of serum IgG antibodies against the SARS-CoV-2 nucleocapsid protein was carried out in the Architect analyser using Abbott's SARS-CoV-2 IgG assay (Abbott, Abbott Park, IL, USA) following manufacturer's instructions. The assay is based on a chemiluminescent microparticle immunoassay and determinations were considered negative or positive depending if

results were < 1.4 or ≥ 1.4 , respectively (cut-off index value) [6]. All samples were processed in the Microbiology and Infectious Disease Department at the Hospital General Universitario Gregorio Marañón.

Microbiological status at the time of sampling was defined as active infection (positive nasal PCR \pm presence of antibodies), past infection (negative nasal PCR + presence of antibodies), or naive participants (negative nasal PCR + absence of antibodies); participants with active or past infections were all together considered infected. Proportions were compared with Epidat v.4.2 (Consellería de Sanidade, Xunta de Galicia, Spain).

Ethics statement. This study was approved by the Ethics Committee of Hospital Gregorio Marañón (CEim; MICRO.HGUGM.2020-019).




Figure 1 Percentage of residents (1A) and staff workers (1B) at each nursing home grouped as either having active infection (positive PCR \pm IgGs), past and cured infection (negative PCR and positive IgGs), and naive patients (negative PCR and negative IgGs). Dates of sample collection are also shown.




Figure 2 Percentage of symptomatic and asymptomatic residents and staff workers, grouped as being actively infected (positive PCR ± IgGs), past and cured infection (negative PCR and positive IgGs), and naïve patients (negative PCR and negative IgGs) at the sampling.

RESULTS

At sampling, 942 surviving residents and 846 staff members were assessed. Median age of the residents was 89 years (IQR: 83.7-92); 77% were female; in the case of staff workers, median age was 46 years (IQR: 38-55); 82.6% were female. The number of residents/staff workers in each nursing home was as follows: NH1 (n=50/44), NH2 (n=284/266), NH3 (n=40/40), NH4 (n=91/121), NH5 (n=129/99), NH6 (n=60/49), NH7 (n=164/116), NH8 (n=55/51), and NH9 (n=64/60). Overall, 224 (23.7%) residents and 127 (15.1%) staff members resulted positive for SARS-CoV-2 by PCR, whereas IgG determination was positive in 665 (70.6%) residents and 398 (47.0%) staff members.

Considering both laboratory results we were able to classify residents as either having active infection (n=224; 23.8%), past infection (n=462; 49.1%), or naïve (n=256; 27.1%); regarding staff members, 127 (15.1%) had active infection, 290 (34.2%) past infection, and 429 were naïve (50.7%). Overall, the percentage of infected participants was significantly higher in residents than in staff members (72.8% vs 49.2%; $P=0.001$). Remarkable differences in terms of infected residents among nursing homes were found (residents among 35.6% and 92.0%; staff workers among 30.0% to 69.4%; Figure 1A and 1B). As the first pandemic wave progressed, the number of affected residents increased, a pattern that was not as clear for staff workers (Figure 1).

The clinical situation at sampling for the 1,269 participants (data was unavailable in 519 participants), residents and staff workers, respectively, was as follows: acute manifestations compatible with COVID-19 (7.3% vs 3.9%; $P<0.01$)

and no manifestations of infection (92.7% vs 96.0%; $P<0.01$). Notably, a large proportion of both asymptomatic and symptomatic residents (69.4% vs 86.6%; $P=0.015$) had PCR positive results (mostly alongside positive IgG determinations) (Figure 2). Out of the naïve resident participants (n=174), only two had had proven COVID-19 and four had had probable COVID-19 (clinically suspected but not demonstrated microbiology); regarding naïve staff members (n=325) three had had proven COVID-19 and 15 had had probable COVID-19.

DISCUSSION

COVID-19 hardly hit Spain during the first wave of the pandemic starting in late February 2020. It had a profound and uneven impact in residents in nursing homes in Madrid, the epicentre of the pandemic in Spain at that time [7]. These institutions represent a setting at a high risk of COVID-19 transmission due to the advanced age of residents and their underlying conditions. SARS-CoV-2 spreading has been deeply described in nursing homes around the world, including other Spanish regions [8], other European countries [9-13], China [14] or the USA [15].

PCR testing for nursing homes residents was implemented in mid-April 2020, which makes impossible to assess COVID-19 attributable mortality before that time. Some estimates suggest that 87% of COVID-19 attributable deaths in Spain occurred among individuals aged 70 years old and above; during the first wave, 13% of all residents died from COVID-19 in Spain [16,17], such a figure rises to 22% in residents over the age of 80 years. In Madrid, 18% of nursing homes residents died from March to May 2020 [18].

Despite the strict restrictions taken during the lockdown, three quarters of the surviving residents in nursing homes in Madrid had some evidence of past or active COVID-19 disease. In contrast, the disease affected staff workers to a lesser extent. As the first wave of cases progressed, the proportion of residents affected also increased. Specific aspects of nursing homes (shared rooms or bathrooms, physically or cognitively impaired residents requiring high-demand care, rotating staff working in several facilities) may have facilitated the rapid spread of viral infections. Unfortunately, restriction policies for visitors in nursing homes implemented as part of the state of emergency declared on March 14 were insufficient to halt further transmissions [19]. The percentage of residents and staff workers who were asymptomatic at the time of sampling illustrates how insufficient the clinical presentation of the disease to control outbreaks resulted, since those asymptomatic cases could have had an important role in transmission [20].

Our study has limitations. First, not all the nursing homes were assessed at the same sampling time; second, clinical situation were not available for all participants; third, clinical situation was recorded at the sample time point and not on previous days. Finally, some nursing homes characteristics such as the proportion of care staff/residents ratio, proportion of shared rooms, among others, was not provided due to the overwhelming situation during the first wave.

In conclusion, COVID-19 affected three quarters of the surviving residents in nursing homes in Madrid, showing how devastating COVID-19 was in such facilities. The high impact suffered in these settings, despite the strict restrictions adopted during the lockdown, demonstrates the ability of SARS-CoV-2 to cause outbreaks.

ACKNOWLEDGMENTS

The authors are grateful to Dainora Jaloveckas for editing assistance. Part of this study was presented at the 31 European Congress of Clinical Microbiology and Infectious Diseases (EC-CMID; Abstract 3078), 2021.

FUNDING

PE (CPII20/00015) is a recipient of a Miguel Servet contract supported by the FIS. JG is a steady researcher contracted by Fundación para Investigación Sanitaria del Hospital Gregorio Marañón.

CONFLICT OF INTEREST

All authors declare no conflicts of interest.

REFERENCES

- World Health Organization. (2020). Coronavirus disease 2019 (COVID-19): Weekly epidemiological update. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200907-weekly-epi-update-4.pdf?sfvrsn=f5f607ee_2.
- Trevisan C, Del Signore S, Fumagalli S, Gareri P, Malara A, Mossello E, et al. Assessing the impact of COVID-19 on the health of geriatric patients: The European GeroCovid Observational Study. *Eur J Intern Med.* 2021; 87:29-35. DOI: 10.1016/j.ejim.2021.01.017.
- Zalakain JD, V. & Suárez-González, A. The COVID-19 on users of Long-Term Care services in Spain. LTCcovid, International Long-Term Care Policy Network, CPEC-LSE, 28 May 2020. 2020.
- Trabucchi M, De Leo D. Nursing homes or besieged castles: COVID-19 in northern Italy. *Lancet Psychiatry.* 2020;7(5):387-8. DOI: 10.1016/S2215-0366(20)30149-8.
- Bouza E, Perez-Granda MJ, Escribano P, Fernandez-Del-Rey R, Pastor I, Moure Z, et al. Outbreak of COVID-19 in a nursing home in Madrid. *J Infect.* 2020;81(4):647-79. DOI: 10.1016/j.jinf.2020.06.055.
- Escribano P, Alvarez-Uria A, Alonso R, Catalan P, Alcalá L, Muñoz P, et al. Detection of SARS-CoV-2 antibodies is insufficient for the diagnosis of active or cured COVID-19. *Sci Rep.* 2020;10(1):19893. DOI: 10.1038/s41598-020-76914-5.
- Soriano V, Ganado-Pinilla P, Sanchez-Santos M, Gomez-Gallego F, Barreiro P, de Mendoza C, et al. Main differences between the first and second waves of COVID-19 in Madrid, Spain. *Int J Infect Dis.* 2021;105:374-6. DOI: 10.1016/j.ijid.2021.02.115.
- Meis-Pinheiro U, Lopez-Segui F, Walsh S, Ussi A, Santaeugenia S, Garcia-Navarro JA, et al. Clinical characteristics of COVID-19 in older adults. A retrospective study in long-term nursing homes in Catalonia. *PLoS One.* 2021;16(7):e0255141. DOI: 10.1371/journal.pone.0255141.
- Araujo O, Sequeira C, Ferre-Grau C, Sousa L. Nursing homes in Portugal during the COVID-19 outbreak: challenges for the future. *Enferm Clin.* 2021. DOI: 10.1016/j.enfcli.2021.09.011.
- Rabilloud M, Riche B, Etard JF, Elsensohn MH, Voirin N, Benet T, et al. COVID-19 outbreaks in nursing homes: A strong link with the coronavirus spread in the surrounding population, France, March to July 2020. *PLoS One.* 2022;17(1):e0261756. DOI: 10.1371/journal.pone.0261756.
- Malara A, Noale M, Abbatecola AM, Borselli G, Cafariello C, Fumagalli S, et al. Clinical Features of SARS-CoV-2 Infection in Italian Long-Term Care Facilities: GeroCovid LTCFs Observational Study. *J Am Med Dir Assoc.* 2022;23(1):15-8. DOI: 10.1016/j.jamda.2021.10.009.
- Roth A, Feller S, Ruhnau A, Plamp L, Viereck U, Weber K, et al. Characterization of COVID-19 outbreaks in three nursing homes during the first wave in Berlin, Germany. *Sci Rep.* 2021;11(1):24441. DOI: 10.1038/s41598-021-04115-9.
- Schultze A, Nightingale E, Evans D, Hulme W, Rosello A, Bates C, et al. Mortality among Care Home Residents in England during the first and second waves of the COVID-19 pandemic: an observational study of 4.3 million adults over the age of 65. *Lancet Reg Health Eur.* 2022;14:100295. DOI: 10.1016/j.lanepe.2021.100295.
- Guo Y, Liu X, Deng M, Liu P, Li F, Xie N, et al. Epidemiology of COVID-19 in older persons, Wuhan, China. *Age Ageing.* 2020;49(5):706-12. DOI: 10.1093/ageing/afaa145.

- 15 McMichael TM, Currie DW, Clark S, Pogojans S, Kay M, Schwartz NG, et al. Epidemiology of Covid-19 in a Long-Term Care Facility in King County, Washington. *N Engl J Med.* 2020;382(21):2005-11. DOI: 10.1056/NEJMoa2005412.
- 16 Spanish National Network for Epidemiological Monitoring (2020a). <https://eng.isciii.es/eng.isciii.es/QueHacemos/Servicios/VigilanciaSaludPublicaRENAV/Paginas/default.html>.
- 17 Spanish National Network for Epidemiological Monitoring (MoMo). (2020b). Data of mortality from March to May 16, 2019. <https://www.isciii.es/QueHacemos/Servicios/VigilanciaSaludPublicaRENAV/EnfermedadesTransmisibles/MoMo/Paginas/MoMo.aspx>, and 2020 https://momo.isciii.es/public/momo/dashboard/momo_dashboard.html#datos.
- 18 Costa-Font J, Jimenez Martin S, Viola A. Fatal Underfunding? Explaining COVID-19 Mortality in Spanish Nursing Homes. *J Aging Health.* 2021;33(7-8):607-17. DOI: 10.1177/08982643211003794.
- 19 Ministerio de la Presidencia relaciones con las cortes y memoria democrática. Royal Decree 463/2020 (March 14) declaring the alarm state for the health crisis management situation caused by COVID-19. BOE no 67 from March 14, 2020. 2020 Mar 14; 67(l): 25390-400. https://www.boe.es/diario_boe/txt.php?id=BOE-A-2020-3692.
- 20 He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med.* 2020;26(5):672-5. DOI: 10.1038/s41591-020-0869-5.